## Is it Flu or Ebola?



## Flu (influenza)



The **flu** is a common contagious respiratory illness caused by flu viruses. The flu is different from a cold.

Flu can cause mild to severe illness, and complications can lead to death.

#### **Ebola**



**Ebola** is a rare and deadly disease caused by infection with an Ebola virus. Sporadic outbreaks have occurred in some African countries since 1976.


## **How Flu Germs Are Spread**





The flu is spread mainly by droplets made when people who have flu cough, sneeze, or talk. Viruses can also spread on surfaces, but this is less common.

People with flu can spread the virus before and during their illness.

## **How Ebola Germs are Spread**







- Infected fruit bats and primates (apes and monkeys)
- And, possibly from contact with semen from a man who has recovered from Ebola (for example, by having oral, vaginal, or anal sex)

## Who Gets The Flu?




Anyone can get the flu.

Some people—like very young children, older adults, and people with some health conditions—are at high risk of serious complications.

# Who Gets Ebola? People most at risk of getting Ebola are







Friends and family who have had unprotected direct contact with blood or body fluids of a person sick with Ebola.

• People with a travel history to countries

#### Signs and Symptoms of Flu



The signs and symptoms of flu usually develop within 2 days after exposure. Symptoms come on quickly and all at once.

## Signs and Symptoms of Ebola



The signs and symptoms of Ebola can appear 2 to 21 days after exposure. The average time is 8 to 10 days. Symptoms of Ebola develop over several days and become progressively more severe.

People with Ebola cannot spread the virus until symptoms appear.



- · Fever or feeling feverish
- Headache
- · Muscle or body aches
- Feeling very tired (fatigue)
- Courdh
- Sore throat
- Runny or stuffy nose



- Fever
- Severe headache
- Muscle pain
- Feeling very tired (fatigue)
- Vomiting and diarrhea develop after 3–6 days
- Weakness (can be severe)
- Stomach pain
- Unexplained bleeding or bruising

For more information about the flu and Ebola, visit www.cdc.gov/flu and www.cdc.gov/ebola.

May 4, 201

CS 252 296-1

Every effort has been made to ensure that the information and the drug names and doses quoted in this Journal are correct. However readers are advised to check information and doses before making prescriptions. Unless otherwise stated the doses quoted are for adults.