

Atypical presentation of COVID-19 in a diabetic patient with malaria-like symptoms: case report

Abrar Bakry Elmalik,¹ Mohamed Eltayieb Elawad,² Alaa Eltayieb Elawad Ahmed³ and Ibrahim A. Ali⁴

1. Medical student, Faculty of Medicine, Omdurman Islamic University, Khartoum, Sudan
2. Medical student, Faculty of Medicine, National Ribat University, Khartoum, Sudan
3. Faculty of Nursing, Sharjah University, Sharjah, United Arab Emirates
4. Assistant Professor of Physiology, Faculty of Medicine, The National Ribat University, Khartoum, Sudan

Correspondence:

Ibrahim Abdelrhim Ali
hemamedicine@gmail.com

Submitted: July 2020

Accepted: August 2020

Published: December 2020

Citation:

Elmalik et al. Atypical Presentation of COVID-19 in Diabetic Patient, with Malaria-like Symptoms: case report. South Sudan Medical Journal 2020; 13(5):196-199 © 2020 The Author (s)
 License: This is an open access article under [CC BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Abstract

Since the emergence of the coronavirus (COVID-19) pandemic beginning in China in 2019, all health workers have faced a difficult challenge. One challenge is a clinical picture in some cases, where the presenting symptoms do not fit with the usual pattern. In this report, we present a case whose initial symptoms were headache, dizziness, and vomiting suggesting malaria bearing in mind that the patient was from Sudan, where there is a high incidence of malaria.

These unusual symptoms may be attributed to the presence of the receptor ACE2, in various body systems. The mild pathological course may be related to the patient's age and the fact that diabetic patients are less likely to develop lung injury, hence the absence of respiratory symptoms.

We recommend that physicians are alert to the variety of different presenting features of COVID-19 and test with Polymerase Chain Reaction (PCR) whenever there is doubt.

Introduction

COVID-19 is a viral disease that infects the respiratory system, and usually, a patient suffers from high fever, fatigue, headache, cough, and shortness of breath. ^[1] This pandemic carries a staggering burden in terms of expenditure on detection, prevention, and treatment, in addition to the loss of life and collapse of the health systems. ^[2] However, many reports have recorded unusual clinical features such as diarrhoea, abdominal pain, testicular pain, and stroke-like symptoms. ^[3,4]

The unusual clinical features cause confusion to the unwary clinician with delayed diagnosis and poorer outcomes. This is especially so if the clinical features are similar to a common disease in the region such as malaria in Sudan, where, despite the lack of statistical studies incidence is estimated at about nine million cases, and the mortality rate about three million cases a year. Thus, malaria is an endemic disease in Sudan. ^[5]

This report presents a case of COVID-19 infection with unusual symptoms and hence emphasises the need to add these features to those when COVID-19 must be considered.

Case Report

A 23-year old female living in Sudan complained of three days of headache and dizziness and vomiting for one day before hospital admission. The moderate bitemporal headache was of sudden onset with a feeling of pressure over the left eye. A day prior to admission, she felt suddenly dizzy that was worse when she lay down and improved on standing, and associated with vomiting after eating. She has type-1 diabetes and is on an insulin pump. She said that she had been in hospital recently with symptoms of hypoglycaemia. A blood sugar level was 50 mg/dl (Reference range 80 – 140 mg/dl). She received 50% dextrose intravenously and was discharged on the same day. The patient told us that she had not complained of headache, dizziness, and vomiting with those characteristics before. Nor did those symptoms appear in her family. She also denied that mosquitoes were in her area, distant travel, or recent direct contact

Table 1. Changes in symptoms since the day of COVID-19 confirmation (Day 8 of the initial admission)

The evening of the first day	Dizziness, cough and severity of the sore throat increases as the moderate headache and repeated sneezing persist
Day 2 and 3	Fatigue and fever intensity increase, previous symptoms persist
Day 4	Fever gradually improving and shortness of breath is emerging again
Day 5	A severe headache reappeared, dizziness and vomiting improved
Day 6 to 8	Mild fever with a less severe headache
Day 9	The disappearance of fatigue and improvement of the general situation but dry cough and sore throat continue
Day 10	The dry cough became productive and sore throat began to improve
Day 11 to 13	Continued productive cough
Day 14	Improvement of productive cough began
Day 15	Mild fever reappears

with a COVID-19 patient.

On examination her pulse rate was 117 beats per minute, respiration rate 17 breaths per minute, temperature 36.9 °C, blood pressure 110/70 mmHg, oxygen saturation 98%. Examination of the nervous system was normal (GCS 15), the cardiovascular system was normal, chest auscultation was clear. However, there was tenderness in the epigastric and periumbilical regions. A peripheral blood film (PBF) for malarial parasites was negative. Other tests showed:

White blood cells (WBCs) was 9×10^9 /L (Reference range $4.5 - 11 \times 10^9$ /L)

Lymphocyte was 1.2×10^9 /L (Reference range $1.5 - 4.5 \times 10^9$ /L)

Monocyte was 0.3×10^9 /L (Reference range $0.2 - 0.8 \times 10^9$ /L)

Eosinophils was 0.2×10^9 /L (Reference range $0 - 0.4 \times 10^9$ /L)

CRP was 1.2 mg/dL (Reference range $0 - 1$ mg/dL)

ESR was 23 mm in one hour (Reference range $1 - 20$ mm in one hour)

In spite of the negative slide for malarial parasites she was treated empirically for malaria with Artemether-Lumefantrine (80/480) tablets, twice daily for three days.

After seven days from admission, the dizziness and vomiting improved. However, the patient began to complain of a mild fever but less severe headache. The next morning, she complained of a dry cough, sore throat, shortness of breath, and repeated sneezing and COVID-19 was suspected. A nasopharyngeal swab was taken and a PCR test was positive for COVID-19 infection.

The patient was isolated. Paracetamol 500 mg tablets were given as required., Vitamin-C one tablet daily to reduce

the damage extent that COVID-19 cause, Vitamin-D capsules 400 IU per day to enhance the immunological function, azithromycin 500 mg on day 1, then 250 mg on days 2-5 due to its antiviral effect and immunity boost, and dextromethorphan hydrobromide was given to ease cough. It was also recommended that an oxygen cylinder be brought for use on demand.

The patient was closely monitored and adequate nourishment was ensured. The changes in symptoms are summarised in Table 1.

Discussion

This case draws attention to the fact that COVID-19 may present as a mild to moderate illness free from complications. Later it may develop into a more serious condition. COVID-19 has been classified into the mild, severe and critical categories listed in Table 2.^[6]

In another study, diabetic patients appeared to be less susceptible to acute lung injury and the development of acute respiratory distress syndrome (ARDS).^[7, 8]

Respiratory complications are among the most serious complications with a high mortality. The age of a patient and severity of the disease are directly related.^[9]

The delay in the diagnosis of COVID-19 because of the unusual initial symptoms exposed doctors and healthcare workers and family members to an increased risk of infection especially since she had visited the hospital twice before the correct diagnosis was made. This is of course in addition to her probable infectivity during a preceding asymptomatic period.

Fever is the most common symptom, followed by cough then fatigue.^[10] But it is now realised there are a variety of other symptoms about which the clinician must be aware. The basis for this variation in presentations may be related to ACE2, as several studies have mentioned. This receptor

Table 2. COVID-19 classification according to severity^[6]

Mild	Absent or mild pneumonia.
Severe	Shortness of breath, respiratory rate more than 29/min, oxygen saturation less than 94%, lung infiltrate 50% within 24 to 48 hours, or PaO ₂ /FiO ₂ ratio less than 300.
Critical	Respiratory failure, multiple organ failure, or septic shock.

is responsible for the formation of vasodilator peptides and found in many places in the human body, including the digestive system and brain hence possibly causing vomiting, headache, and dizziness.^[11,12]

Consideration of the differential diagnoses is crucial when presented with non-specific symptoms that might be caused by a number of infectious diseases, especially if they are common in the region. In Thailand, which is a tropical area, it is likely that COVID-19 will be confused with tropical diseases. A series of cases (48) were reported with a petechial rash, and thrombocytopenia which were initially thought to indicate dengue fever.^[13]

In our case the main complaints in the first presentation were headache, dizziness, and vomiting, symptoms that could indicate malaria, neurological disease, or meningitis. However, the neurological examination was normal. Although the peripheral blood film did not show malarial parasites, it is clinical practice in Sudan to consider a diagnose malaria if eosinophils are more than or equal to 0.2, and there are increased monocytes. However, the normal WBCs, low lymphocyte count, and increased ESR and CRP indicate a coronavirus.^[14,15]

Conclusion

This case adds to our knowledge about the presentations of COVID-19 and how they might mimic other conditions, malaria in particular. Some differentials of the WBCs such as high monocyte and eosinophils counts suggested a diagnosis of malaria.

We recommend that clinicians should be alert to the possibility of COVID-19 whenever a patient presents with non-specific features and make modifications to the triage system to include recorded unusual symptoms. If there is any doubt a test for COVID-19 should be requested especially if the patient has another condition (e.g. diabetes, chronic respiratory disease) that puts them into a higher risk group that should be managed by intensive care. A delayed diagnosis may increase the severity of the disease, resulting in complications and even death. In addition, the risk of further spread of infection is increased.

References

1. Tian S, Hu N, Lou J, et al. Characteristics of COVID-19 infection in Beijing. *Journal of Infection* April 2020;80(4):401-406.

2. Livingston E, Bucher K. Coronavirus disease 2019 (COVID-19) in Italy. *JAMA* 2020;323(14):1335-1335.
3. Kim J, Thomsen T, Sell N, Goldsmith AJ. Abdominal and testicular pain: An atypical presentation of COVID-19. *American Journal of Emergency Medicine* July 2020;38(7):1542.E1-1542.E3
4. Avula A, Nalleballe K, Narula N. et al. COVID-19 presenting as stroke. *Brain, behavior, and immunity* 2020;87:115-119.
5. Abdalla SI, Malik EM, Ali KM. The burden of malaria in Sudan: incidence, mortality and disability-adjusted life-years. *Malaria Journal* 2007 Dec 1;6(1):97.
6. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020 Apr 7;323(13):1239-42.
7. Honiden S, Gong MN. Diabetes, insulin, and development of acute lung injury. *Critical Care Medicine* 2009;37(8): 2455.
8. Yu S, Christiani DC, Thompson BT et al. Role of diabetes in the development of acute respiratory distress syndrome. *Critical Care Medicine* 2013;41(12):2720.
9. Onder G, Rezza G, Brusaferro S. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *JAMA* 2020;323(18):1775-1776.
10. Wang D, Hu B, Hu C et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323(11):1061-1069.
11. Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Frontiers of Medicine*. 2020 May 9;14(2):185-92.

12. Doobay M, Talman LS, Obr TD. et al. Differential expression of neuronal ACE2 in transgenic mice with overexpression of the brain renin-angiotensin system. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 2007; 292(1):R373-R381.
13. Joob B, Wiwanitkit V. COVID-19 can present with a rash and be mistaken for Dengue. *Journal of the American Academy of Dermatology*. 2020 May;82(5):e177.
14. Elsayid M, Elbasheer MA, Othman AA, Belal RH, Elfaki TE. Estimation of complete blood count and platelets indices in Sudanese patients with malaria. *American Journal of Research Communication*. 2017;5(1):29-38.
15. Fan BE, Chong VC, Chan SS, Lim GH, Lim KG, Tan GB, Mucheli SS, Kuperan P, Ong KH. Hematologic parameters in patients with COVID-19 infection. *American Journal of Hematology*. 2020 Jun;95(6):E131-4.